

AS-ITP-98-12, hep-ph/9810491; To appear in Phys. Rev. D

SO(3) Gauge Symmetry and Neutrino-Lepton Flavor Physics

Yue-Liang Wu

*Institute of Theoretical Physics, Academia Sinica,
P.O. Box 2735, Beijing 100080, P.R. China*

Abstract

Based on the SO(3) gauge symmetry for three family leptons and general see-saw mechanism, we present a simple scheme that allows three nearly degenerate Majorana neutrino masses needed for hot dark matter. The vacuum structure of the spontaneous SO(3) symmetry breaking can automatically lead to a maximal CP-violating phase. Thus the current neutrino data on both the atmospheric neutrino anomaly and solar neutrino deficit can be accounted for via maximal mixings without conflict with the current data on the neutrinoless double beta decay. The model also allows rich interesting phenomena on lepton flavor violations.

PACS numbers: 12.15F, 11.30H

Typeset using REVTeX

Studies on neutrino physics have resulted in the following observations: i), The Super-Kamiokande data [1] on atmospheric neutrino anomaly provide a strong evidence that neutrinos are massive; ii), The current Super-Kamiokande data on solar neutrino [2] cannot decisively establish whether the solar neutrino deficit results from ‘just so’ oscillations [3] or MSW solutions [4] with small/large mixing angles [5]. iii), To describe all the neutrino phenomena such as the atmospheric neutrino anomaly, the solar neutrino deficit and the results from the LSND experiment, it is necessary to introduce a sterile neutrino [6]. It indicates that with only three light neutrinos, one of the experimental data must be modified; iv), The current experimental data cannot establish whether neutrinos are Dirac-type or Majorana-type. The failure of detecting neutrinoless double beta decay only provides, for Majorana-type neutrinos, an upper bound on an ‘effective’ electron neutrino mass; v), Large neutrino masses around several electron volts may play an important role in the evolution of the large-scale structure of the universe [7]. Having noticed these facts, we will present in this paper a conservative consideration in which only three light neutrinos are involved.

To consider massive neutrinos, it is necessary to extend the minimal standard model (SM). The greatest success of the SM is its gauge symmetry structure $SU_L(2) \times U_Y(1)$. As a simple extension of the standard model, it is of interest to introduce a flavor symmetry among the three families of the leptons. In this paper, we will choose the $SO(3)$ symmetry group to describe the lepton flavors and treat it as a gauge symmetry. As usual, we will also introduce an additional $U(1)$ symmetry to construct the needed interaction terms. Let us begin with the following $SO(3) \times SU(2)_L \times U(1)_Y$ invariant lagrangian for leptons

$$\begin{aligned} \mathcal{L} = & \mathcal{L}_{YM} + \bar{L}_i i\gamma^\mu D_\mu L_i + \bar{e}_R i\gamma^\mu D_\mu e_{R\ i} + \bar{N}_R i\gamma^\mu D_\mu N_{R\ i} + \bar{V} i D_\mu V + \bar{E} i\gamma^\mu D_\mu E \\ & + y_1 \bar{L}_i \varphi_i V + y_2 \bar{V} \phi_1 E + y_3 \bar{E} \varphi_i e_{R\ i} + y_0 \bar{L}_i \phi_2 N_{R\ i} + S(x) \bar{N}_{R\ i} N_{R\ i}^c + H.c. \\ & + M_1 \bar{E} E + M_2 \bar{V} V + D^\mu \phi_1^* D_\mu \phi_1 + D^\mu \phi_2^* D_\mu \phi_2 + D^\mu \varphi^* D_\mu \varphi - V(\varphi_i, \phi_1, \phi_2, S) \end{aligned} \quad (1)$$

with

$$\begin{aligned} D_\mu L_i &= (\partial_\mu + ig_1 \frac{1}{2} B_\mu - ig_2 W_\mu^k \frac{\tau^k}{2}) L_i - ig_3' A_\mu^k \frac{(t^k)_{ij}}{2} L_j \\ D_\mu e_{R\ i} &= (\partial_\mu + ig_1 B_\mu) e_{R\ i} - ig_3' A_\mu^k \frac{(t^k)_{ij}}{2} e_{R\ j} \\ D_\mu N_{R\ i} &= \partial_\mu N_{R\ i} - ig_3' A_\mu^k \frac{(t^k)_{ij}}{2} N_{R\ j} \\ D_\mu V &= (\partial_\mu - ig_1 \frac{1}{2} B_\mu - ig_2 W_\mu^i \frac{\tau^i}{2}) V \\ D_\mu E &= (\partial_\mu + ig_1 B_\mu) E \\ D_\mu \varphi_i &= \partial_\mu \varphi_i - ig_3' A_\mu^k \frac{(t^k)_{ij}}{2} \varphi_j \\ D_\mu \phi_1 &= (\partial_\mu - ig_1 \frac{1}{2} B_\mu - ig_2 W_\mu^k \frac{\tau^k}{2}) \phi_1 \\ D_\mu \phi_2 &= (\partial_\mu + ig_1 \frac{1}{2} B_\mu - ig_2 W_\mu^k \frac{\tau^k}{2}) \phi_2 \end{aligned} \quad (2)$$

Where $i, j = 1, 2, 3$ being family indices and t^k ($k = 1, 2, 3$) are the three $SO(3)$ generators. The $U(1)$ charges of the various fields are chosen as: $(\varphi_i(x), V(x), E(x), N_{R\ i}(x),$

$S(x)$, $e_{R\ i}(x)$, $L_i(x)$, $\phi_1(x)$, $\phi_2(x)$) = (-1/2, 1/2, 1/2, -1, -2, 1, 0, 0, 1), so that the theory is anomaly free. $\bar{L}_i(x) = (\bar{\nu}_i, \bar{e}_i)_L$ are the $SU(2)_L$ doublet leptons. $e_{R\ i}$ and $\nu_{R\ i}$ are right-handed charged leptons and neutrinos. $\phi_1(x)$ and $\phi_2(x)$ are the two Higgs doublets. $\varphi^T = (\varphi_1(x), \varphi_2(x), \varphi_3(x))$ is a complex $SO(3)$ triplet scalar and $S(x)$ is a complex singlet scalar. $V(x)$ is an $SU(2)_L$ doublet vector-like lepton with mass M_2 and $E(x)$ is a singlet vector-like electron with mass M_1 . The three Majorana neutrinos $N_{R\ i}$ also belong to the $SO(3)$ triplet representation. $A_\mu = A_\mu^k(x)t^k/2$ is the $SO(3)$ gauge field and g'_3 is the $SO(3)$ gauge coupling constant. $\mathcal{L}_{YM}$ represents the pure Yang-Mills gauge interactions and $V(\varphi_i, \phi_1, \phi_2, S)$ denotes the Higgs potential. If the singlet scalar $S(x)$ carries two unit lepton number, the lepton number is also conserved in the above lagrangian. The lepton number and $U(1)$ symmetries are supposed to be spontaneously broken down at very high energy scales $< S(x) > = M_N \simeq 10^{13}\text{GeV}$ and their effects will be not discussed in this paper. The mass scales M_1 and M_2 are assumed to be above the energy scales of the $SO(3)$ flavor symmetry breaking which could occur at low energy scales. Adopting the general see-saw mechanism, we obtain the following effective lagrangian at the leading order

$$\mathcal{L} = C_1 \frac{\varphi_i \varphi_j}{M_1 M_2} \bar{L}_i \phi_1 e_{R\ j} + C_0 \frac{1}{M_N} \bar{L}_i \phi_2 \phi_2^T L_i^c \quad (3)$$

with $C_1 = y_1 y_2 y_3$ and $C_0 = y_0^2$. This effective lagrangian remains invariant under $SO(3) \times SU_L(2) \times U_Y(1)$ action. After the symmetry is broken down to $U_{em}(1)$, one yields the following simple mass matrices for the Majorana neutrinos and charged leptons

$$M_\nu = m_0 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad (4)$$

and

$$M_e = \frac{m_1}{\sigma^2} \begin{pmatrix} \hat{\sigma}_1^2 & \hat{\sigma}_1 \hat{\sigma}_2 & \hat{\sigma}_1 \hat{\sigma}_3 \\ \hat{\sigma}_1 \hat{\sigma}_2 & \hat{\sigma}_2^2 & \hat{\sigma}_2 \hat{\sigma}_3 \\ \hat{\sigma}_1 \hat{\sigma}_3 & \hat{\sigma}_2 \hat{\sigma}_3 & \hat{\sigma}_3^2 \end{pmatrix} \quad (5)$$

with $m_0 = C_0 v_2^2 / M_N$ and $m_1 = C_1 v_1 \sigma^2 / M_1 M_2$. Where $v_{1,2} = \langle \phi_{1,2} \rangle$, $\hat{\sigma}_i = \langle \varphi_i \rangle$ and $\sigma = \sqrt{|\hat{\sigma}_1|^2 + |\hat{\sigma}_2|^2 + |\hat{\sigma}_3|^2}$ are the vacuum expectation values of the corresponding fields after spontaneous symmtry breaking. Here the values of the quantities $y_2 v_1$ and $|\hat{\sigma}_i|$ are considered to be smaller than the mass scales M_1 and M_2 , i.e., $y_2 v_1, |\hat{\sigma}_i| < M_1, M_2$.

As the $SO(3)$ flavor symmetry is treated as a gauge symmetry, one can always express the complex $SO(3)$ triplet scalar fields $\varphi_i(x)$ in terms of the three rotational fields and three amplitude fields. This is analogous to $SU(2)$ gauge symmetry, the complex $SU(2)$ doublet scalar field can always be expressed in terms of three $SU(2)$ 'rotational' fields and one amplitude field. As the $SO(3)$ rotation matrix is real, which is unlike to the $SU(2)$ rotation matrix that is complex, one of the three amplitude fields of the complex $SO(3)$ triplet scalar must be a pure imaginary field so that one can generate the complex $SO(3)$ triplet scalar fields $\varphi_i(x)$ by an $SO(3)$ field $O(x) = e^{i\eta_i(x)t^i} \in SO(3)$ action on the three amplitude fields. Explicitly, one has

$$\begin{pmatrix} \varphi_1(x) \\ \varphi_2(x) \\ \varphi_3(x) \end{pmatrix} = e^{i\eta_i(x)t^i} \frac{1}{\sqrt{2}} \begin{pmatrix} \rho_1(x) \\ i\rho_2(x) \\ \rho_3(x) \end{pmatrix} \quad (6)$$

where the three real rotational fields $\eta_i(x)$ and the three real amplitude fields $\rho_i(x)$ reparameterize the six real fields of the complex triplet scalar field $\varphi(x)$. In general there are three possible assignments for the imaginary amplitude. In this note, we only consider the case that the imaginary part is assigned to the second amplitude field. From the Higgs mechanism, the three rotational fields $\eta_i(x)$ will be ‘eaten’ by the three gauge fields $A_\mu^i(x)$ after spontaneous symmetry breaking

$$\rho_i(x) \rightarrow \sigma_i + \rho_i(x) \quad (7)$$

For this case, one has $\hat{\sigma}_1 = \sigma_1$, $\hat{\sigma}_2 = i\sigma_2$ and $\hat{\sigma}_3 = \sigma_3$. With this analysis, it is seen that when three amplitude fields get nonzero vacuum expectation values, an $SO(3)$ gauge theory with one complex triplet Higgs field always spontaneously breaks CP invariance via a $\pi/2$ phase.

It is easy to check that the matrix M_e is a rank one matrix and diagonalized by a unitary matrix U_e via

$$D_e = U_e^\dagger M_e U_e^* \quad (8)$$

with

$$D_e = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & m_1 \end{pmatrix} \quad (9)$$

and

$$U_e = \begin{pmatrix} -ic_1 & c_2 s_1 & s_1 s_2 \\ -s_1 & ic_1 c_2 & ic_1 s_2 \\ 0 & -s_2 & c_2 \end{pmatrix} \quad (10)$$

where we have used the notations $s_1 \equiv \sin \theta_1 = \sigma_1/\sigma_{12}$, $s_2 \equiv \sin \theta_2 = \sigma_{12}/\sigma$, $\sigma_{12} = \sqrt{\sigma_1^2 + \sigma_2^2}$ and $\sigma = \sqrt{\sigma_{12}^2 + \sigma_3^2}$.

As the neutrinos are in the mass eigenstate basis, the CKM-type lepton mixing matrix U_{LEP} defined in the charged weak gauge interactions

$$\mathcal{L}_W = \bar{e}_L \gamma^\mu U_{LEP} \nu_L W_\mu^- + H.c. \quad (11)$$

is simply given by

$$U_{LEP} = U_e^\dagger = \begin{pmatrix} ic_1 & -s_1 & 0 \\ c_2 s_1 & -ic_1 c_2 & -s_2 \\ s_1 s_2 & -ic_1 s_2 & c_2 \end{pmatrix} \quad (12)$$

Before proceeding, we would like to address the following points: Firstly, as neutrino mass matrix is a Majorana one, the complex U_{LEP} cannot be absorbed. Secondly, as the neutrino

masses are degenerate at the leading order of the present considerations, the unitary matrix U_{LEP} is arbitrary up to multiplication by an arbitrary orthogonal matrix from the right. This arbitrariness can be fixed by eliminating the degeneracy of the neutrino masses, this may be realized by considering additional new contributions and high order corrections to the neutrino masses. It will also be seen below that such an arbitrariness is actually prevented from the $SO(3)$ gauge interactions of the neutrinos. Thirdly, as the matrix element $(U_{LEP})_{13} = 0$ such a unitary matrix can always be transformed into an orthogonal matrix by a phase redefinition of the left-handed neutrinos and charged leptons. For the above case, it can be realized by redefining $e_L \rightarrow ie_L$ and $\nu_{\mu L} \rightarrow i\nu_{\mu L}$, one then has

$$U_{LEP} \rightarrow O_{LEP} = \begin{pmatrix} c_1 & -s_1 & 0 \\ c_2 s_1 & c_1 c_2 & -s_2 \\ s_1 s_2 & c_1 s_2 & c_2 \end{pmatrix} \quad (13)$$

However, we would like to emphasize that such a phase redefinition is not trivial due to Majorana neutrinos and $SO(3)$ gauge interactions. This is because the neutrino mass matrix becomes, after the phase redefinition $\nu_{\mu L} \rightarrow i\nu_{\mu L}$, the following form

$$M_\nu \rightarrow M_\nu = m_0 \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad (14)$$

which is no longer a unit matrix and the neutrinos are not in the physical mass basis. A direct physical effect of such a CP phase can be seen in the processes concerning the neutrinoless double beta decay. One can also explicitly see below that the phase redefinition will lead to a change of the phases of the couplings in the $SO(3)$ gauge interactions of the charged leptons and neutrinos.

When transferring the neutrino mass basis into the weak gauge and charged lepton mass basis, the neutrino mass matrix $M_\nu = m_0 U_{LEP} U_{LEP}^T$ is found to have the following form

$$M_\nu = m_0 \begin{pmatrix} s_1^2 - c_1^2 & 2ic_1 s_1 c_2 & 2ic_1 s_1 s_2 \\ 2ic_1 s_1 c_2 & c_2^2(s_1^2 - c_1^2) + s_2^2 & c_2 s_2(s_1^2 - c_1^2) - c_2 s_2 \\ 2ic_1 s_1 s_2 & c_2 s_2(s_1^2 - c_1^2) - c_2 s_2 & s_2^2(s_1^2 - c_1^2) + c_2^2 \end{pmatrix} \quad (15)$$

which can be diagonalized by a unitary matrix U_ν via $U_\nu^T M_\nu U_\nu$ with $U_\nu = U_{LEP}$.

To describe the realistic world, one needs to consider mechanisms that will split the degeneracy of the three neutrinos and produce the correct muon and electron masses. These mechanisms can arise from high order radiative corrections or suppressed contributions due to additional new interaction terms. As the mass differences among the three neutrinos are expected to be very small, the muon and electron masses are known to be much smaller than the tau mass, it is then reasonable to obtain a realistic pattern which will not significantly differ from the current pattern. For simplicity of discussions, we shall leave such constructions elsewhere [8] and consider the present scheme, similar to some other schemes such as the democratic scheme, to be the first essential step.

As an approximate estimation, the current experimental data [9] from neutrinoless double beta decay indicates that

$$\langle m_{\nu_e} \rangle = m_0(s_1^2 - c_1^2) < 0.46eV \quad (16)$$

The required hot dark matter needs $m_0 \simeq 1.5$ eV. Combinations of these two experimental data will result in a bound on θ_1

$$36^\circ < \theta_1 < 54^\circ, \quad or, \quad \sin^2 2\theta_1 > 0.9 \quad (17)$$

which shows that when neutrinos do play an essential role for the evolution of large scale structure of the universe, the solar neutrino deficit appears to be explained by the $\nu_e - \nu_\mu$ mixing via ‘Just So’ oscillations or MSW solution with large mixing angle. Furthermore, the current Super-Kamiokande data on atmospheric neutrinos require

$$\sin^2 2\theta_2 > 0.8, \quad or, \quad 32^\circ < \theta_2 < 58^\circ \quad (18)$$

These consequences motivate us to consider scenarios in which the two vacuum expectation values of σ_1 and σ_2 are equal, i.e., $\sigma_1 = \sigma_2$ (or $s_1 = 1/\sqrt{2}$), the neutrino mass and mixing matrices can be simply written as

$$M_\nu = m_0 \begin{pmatrix} 0 & ic_2 & is_2 \\ ic_2 & s_2^2 & -c_2 s_2 \\ is_2 & -c_2 s_2 & c_2^2 \end{pmatrix} \quad (19)$$

and

$$U_{LEP} = \begin{pmatrix} \frac{1}{\sqrt{2}}i & -\frac{1}{\sqrt{2}} & 0 \\ \frac{1}{\sqrt{2}}c_2 & -\frac{1}{\sqrt{2}}c_2i & -s_2 \\ \frac{1}{\sqrt{2}}s_2 & -\frac{1}{\sqrt{2}}s_2i & c_2 \end{pmatrix} \quad (20)$$

which arrives at the pattern suggested in [12].

Two particular interesting cases which have been widely discussed in the literature can be easily achieved by further considering the following two cases: First, $\sigma_3^2 = \sigma_1^2 + \sigma_2^2$ and $\sigma_1 = \sigma_2$, namely, $s_1 = s_2 = 1/\sqrt{2}$, one then arrives at the bi-maximal mixings [10,11] with a maximal CP-violating phase. Explicitly, the neutrino mass and mixing matrices read

$$M_\nu = m_0 \begin{pmatrix} 0 & \frac{1}{\sqrt{2}}i & \frac{1}{\sqrt{2}}i \\ \frac{1}{\sqrt{2}}i & \frac{1}{2} & -\frac{1}{2} \\ \frac{1}{\sqrt{2}}i & -\frac{1}{2} & \frac{1}{2} \end{pmatrix} \quad (21)$$

and

$$U_{LEP} = \begin{pmatrix} \frac{1}{\sqrt{2}}i & -\frac{1}{\sqrt{2}} & 0 \\ \frac{1}{2} & -\frac{1}{2}i & -\frac{1}{\sqrt{2}} \\ \frac{1}{2} & -\frac{1}{2}i & \frac{1}{\sqrt{2}} \end{pmatrix} \quad (22)$$

which yields the pattern discussed recently by Georgi and Glashow [13].

Second, when three vacuum expectation values σ_i (i=1,2,3) are democratic, i.e., $\sigma_3 = \sigma_1 = \sigma_2$, namely $s_1 = 1/\sqrt{2}$ and $s_2 = \sqrt{2/3}$, we then obtain the democratic mixing [14,15] with a maximal CP-violating phase. The explicit neutrino mass and mixing matrices are given by

$$M_\nu = m_0 \begin{pmatrix} 0 & \frac{1}{\sqrt{3}}i & \frac{2}{\sqrt{6}}i \\ \frac{1}{\sqrt{3}}i & \frac{2}{3} & -\frac{\sqrt{2}}{3} \\ \frac{2}{\sqrt{6}}i & -\frac{\sqrt{2}}{3} & \frac{1}{3} \end{pmatrix} \quad (23)$$

and

$$U_{LEP} = \begin{pmatrix} \frac{1}{\sqrt{2}}i & -\frac{1}{\sqrt{2}} & 0 \\ \frac{1}{\sqrt{6}} & -\frac{1}{\sqrt{6}}i & -\frac{2}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{3}}i & \frac{1}{\sqrt{3}} \end{pmatrix} \quad (24)$$

which comes to a similar form provided by Mohapatra [16] for the degenerate neutrino scenario.

So far we have mainly focused on the neutrino masses and mixings. In fact, rich new interesting physical phenomena may arise from the SO(3) gauge interactions. We would like to address the following important point that the redefinition of the phases of the leptons is not trivial in the present model, which is unlike to the standard model. This is because the SO(3) gauge fields couple to lepton flavor changing neutral currents. Explicitly, the SO(3) gauge interactions in the mass eigenstate of the leptons have the following form

$$\mathcal{L}_F = \frac{1}{2}g'_3 \bar{\nu}_L t^i \gamma^\mu \nu_L A_\mu^i + \frac{1}{2}g'_3 \bar{e}_L K_e^i \gamma^\mu e_L A_\mu^i - \frac{1}{2}g'_3 \bar{e}_R K_e^{i*} \gamma^\mu e_R A_\mu^i \quad (25)$$

with $K_e^i = U_e^\dagger t^i U_e$. Using the relation $K_e^i = U_e^\dagger t^i U_e = K_e^{ia} t^a$ with t^a ($a = 1, \dots, 9$) being the nine U(3) generators, the above interactions can be reexpressed as

$$\mathcal{L}_F = \frac{1}{2}g'_3 \bar{\nu}_L t^i \gamma^\mu \nu_L A_\mu^i + \frac{1}{2}g'_3 \bar{e} \bar{K}_e^i \gamma^\mu e A_\mu^i + \frac{1}{2}g'_3 \bar{e} \tilde{K}_e^i \gamma^\mu e A_\mu^i \quad (26)$$

with $\bar{K}_e^i = K_e^{ij} t^j$ for vector currents and $\tilde{K}_e^i = K_e^{ia'} t^{a'}$ for axial vector currents. Where t^i ($i = 1, 2, 3$) are the three complex (or antisymmetric) generators of U(3) and $t^{a'}$ ($a' = 4, \dots, 9$) are the six real (or symmetric) generators of U(3). Note that the above interaction forms are given at the SO(3) gauge basis. After spontaneous symmetry breaking of SO(3), the gauge fields A_μ^i receive masses by 'eating' the three rotational fields $\eta^i(x)$. For the SO(3) vacuum structure given above, A_μ^1 and A_μ^3 are not in the mass eigenstates since they mix each other. Denoting the physical gauge fields as F_μ^i , we have $A_\mu^i = O_F^{ij} F_\mu^j$. Explicitly, it reads

$$\begin{pmatrix} A_\mu^1 \\ A_\mu^2 \\ A_\mu^3 \end{pmatrix} = \begin{pmatrix} c_3 & 0 & -s_3 \\ 0 & 1 & 0 \\ s_3 & 0 & c_3 \end{pmatrix} \begin{pmatrix} F_\mu^1 \\ F_\mu^2 \\ F_\mu^3 \end{pmatrix} \quad (27)$$

with the mixing angle $s_3 = \sin \theta_3 = \sigma_1 / \sigma_{13}$ ($\sigma_{13} = \sqrt{\sigma_1^2 + \sigma_3^2}$). Their physical masses are:

$$m_{F_1} = \frac{1}{2}g'_3 \sigma_2, \quad m_{F_2} = \frac{1}{2}g'_3 \sigma_{13}, \quad m_{F_3} = \frac{1}{2}g'_3 \sigma \quad (28)$$

In the physical mass basis of leptons and gauge bosons, the above interactions will be

$$\mathcal{L}_F = \frac{1}{2}g'_3 \bar{\nu}_L t^j O_F^{ji} \gamma^\mu \nu_L F_\mu^i + \frac{1}{2}g'_3 \bar{e}_L V_e^i \gamma^\mu e_L F_\mu^i - \frac{1}{2}g'_3 \bar{e}_R V_e^{i*} \gamma^\mu e_R F_\mu^i \quad (29)$$

with $V_e^i = K_e^j O_F^{ji} = U_e^\dagger t^j U_e O_F^{ji}$. To be explicit, for the case considered above, we have

$$K_e^1 = \begin{pmatrix} 2c_1s_1 & ic_2(s_1^2 - c_1^2) & is_2(s_1^2 - c_1^2) \\ -ic_2(s_1^2 - c_1^2) & 2c_1s_1c_2^2 & 2c_1s_1c_2s_2 \\ -is_2(s_1^2 - c_1^2) & 2c_1s_1c_2s_2 & 2c_1s_1s_2^2 \end{pmatrix} \quad (30)$$

$$K_e^2 = \begin{pmatrix} 0 & c_1s_2 & -c_1c_2 \\ c_1s_2 & 0 & is_1 \\ -c_1c_2 & -is_1 & 0 \end{pmatrix} \quad (31)$$

and

$$K_e^3 = \begin{pmatrix} 0 & is_1s_2 & -is_1c_2 \\ -is_1s_2 & 2c_1c_2s_2 & (s_2^2 - c_2^2)c_1 \\ is_1c_2 & (s_2^2 - c_2^2)c_1 & -2c_1c_2s_2 \end{pmatrix} \quad (32)$$

and

$$\begin{aligned} V_e^1 &= \cos \theta_3 K_e^1 + \sin \theta_3 K_e^3, \\ V_e^2 &= K_e^2, \\ V_e^3 &= -\sin \theta_3 K_e^1 + \cos \theta_3 K_e^3 \end{aligned} \quad (33)$$

If the mixing matrix U_e does not significantly affected when muon lepton gets its physical mass, the current data on lepton flavor violating process $\mu \rightarrow 3e$ with $Br(\mu \rightarrow 3e) < 1 \times 10^{-12}$ [17] lead to the following constraints on the SO(3) symmetry breaking scale

$$\sigma_1 \sim \sigma_2 \sim \sigma_3 > 10^3 v \quad (34)$$

with $v = 246$ GeV being the weak symmetry breaking scale.

In conclusion, it has been shown that the SO(3) gauge symmetry for lepton flavors appears to have some remarkable features which are applicable to the current interesting phenomena concerning neutrinos.

Acknowledgments: This work was supported in part by the NSF of China under the grant No. 19625514.

REFERENCES

- [1] T. Kajita, Talk at XVIII Intern'l Conf. on Neutrino Physics and Astrophysics, Takayama, Japan; Super-Kamiokande Collaboration, Y.Fukuda *et al.*, Phys. Rev. Lett. **81**, 1562 (1998), hep-ex/9807003.
- [2] E.g., Y. Suzuki, Talk at XVIII Intern'l Conf. on Neutrino Physics and Astrophysics, Takayama, Japan; Y.Fukuda *et al.*, Phys. Rev. Lett. **81**, 1158 (1998).
- [3] V. Barger, K. Whisnant and R.J.N Phillips, Phys. Rev. **D24** 538 (1981); S.L. Glashow and L.M. Krauss, Phys. Lett. **B190** (199) (1987).
- [4] L. Wolfenstein, Phys. Rev. **D17** 2369 (1978); S.P. Mikheyev and A. Yu. Smirnov, Sov. J. Nucl. Phys. 42 913 (1986).
- [5] J.N. Bahcall, P. Krastev and A.Yu. Smirnov, Phys. Rev. **D58**, 096016 (1998), hep-ph/9807216.
- [6] D.O. Caldwell and R.N. Mohapatra, Phys. Rev. **D48** 3259 (1993); J.T. Peltoniemi and J.W.F. Valle, Nucl. Phys. B406 409 (1993); R. Foot and R.R. Volkas, Phys. Rev. **D52** 6595 (1995); K.C. Chou and Y.L. Wu, Nucl. Phys. Proc. Suppl. **52A** (1997) 159-163, hep-ph/9610300; hep-ph/9708201; Zhou Guang-Zhao (K.C. Chou) and Wu Yue-Liang (Y.L. WU) Science in China, 39A, 65 (1996), hep-ph/9508402; S.M. Bilenky, C. Giunti and W. Grimus, Eur. Phys. J. **C1** 247 (1998); V. Barger, K. Whisnant and T.J. Weiler, Phys. Lett. **B427**, 97 (1998), hep-ph/9712495; S.C. Gibbons, R.N. Mohapatra, S. Nandi and A. Raychaudhuri, Phys. Lett. **B430**, 296 (1998), hep-ph/9803299; V. Barger, S. Pakvasa, T.J. Weiler and K. Whisnant, Phys. Rev. **D58**, 093016 (1998), hep-ph/9806328; A. S. Joshipura and A. Yu. Smirnov, Phys. Lett. **B439**, 103 (1998), hep-ph/9806376.
- [7] J.R. Primack and M. A. K. Gross, astro-ph/9810204, to be published in the proceedings of the Xth Rencontres de Blois, "The Birth of Galaxies", 28 June - 4 July 1998; J.R. Primack, J. Holtzman, A. Klypin and D.O. Caldwell, Phys. Rev. Lett. **74** 2160 (1995).
- [8] Y.L. Wu, AS-ITP-99-01, hep-ph/9901245; AS-ITP-99-04, hep-ph/9901320.
- [9] Heideberg-Moscow collaboration, M. Günther *et al.*, Phys. Rev. D55, 54 (1997); L. Baudis *et al.*, Phys. Lett. **B407** 219 (1997); Klapdor-Kleingrothaus, hep-ex/9802007.
- [10] 'Bi-maximal mixing matrix' without maximal CP-violating phase was described by V. Barger, S. Pakvasa, T. Weiler and K. Whisnant, Phys. Lett. **B437**, 107 (1998), hep-ph/9806387 ; A.J. Baltz, A.S. Goldhaber and M. Goldhaber, hep-ph/9806540; M. Jezabek and Y. Sumino, hep-ph/9807310.
- [11] Some interesting models with bi-maximal mixing pattern may be found in papers: Y. Nomura and T. Yanagida, hep-ph/9807325; R.N. Mohapatra and S. Nussinov, hep-ph/9809415; M.Fukugita, M. Tanimoto and T. Yanagida, hep-ph/9809554; C. Jarlskog, M. Matsuda, S. Skadhauge and M. Tanimoto, hep-ph/9812282.
- [12] F. Vissani, hep-ph/9708483.
- [13] H. Georgi and S.L. Glashow, HUTP-98/A060, hep-ph/9808293.

- [14] ‘Democratic mixing matrix’ without maximal CP-violating phase was described by H. Fritzsch and Z.Z. Xing, Phys. Lett. **B 372** 265 (1996).
- [15] Some interesting models with democratic mixing pattern may be found in papers:
M.Fukugita, M. Tanimoto and T. Yanagida, Phys. Rev. **D57**, 4429 (1998);
Y. Koide, Phys. Rev. **D 39** 1391 (1989);
R.N. Mohapatra and S. Nussinov, hep-ph/9808301;
M. Tanimoto, hep-ph/9807283.
- [16] R.N. Mohapatra, hep-ph/9808284.
- [17] See summary talk by D. Bryman at ICHEP98, Vancouver, Canada, 1998.